

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DICER 500

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA
present as the dimethylamine salt

GROUP I HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugarcane, turf, rice and non-crop areas as per the Directions for Use.

**IMPORTANT: READ THE LABEL AND ATTACHED
LEAFLET BEFORE OPENING OR USING THIS PRODUCT**

CONTENTS: 5 Litres
APVMA Approval No.: 65687/51582

Hextar

HEXTAR CHEMICALS PTY. LTD ACN No. 134 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008. Telephone: (03) 8687 9509 Fax: (03) 8687 9078

DICER 500

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and setup for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of DICER 500 Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

In a Transport
Emergency Dial
000
Police or Fire Brigade



Hextar

HEXTAR CHEMICALS PTY. LTD. ACN NO.: 114 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008.

Telephone: (03) 8687 9509 Fax: (03) 8687 9078



9555497505640

Batch No.:

Date of Manufacture:

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DICER 500

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA present as the dimethylamine salt

GROUP I HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugarcane, turf, rice and non-crop areas as per the Directions for Use.

IMPORTANT: READ THE LABEL AND ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

CONTENTS: 20 Litres

APVMA Approval No.: 65687/51582

Hexar

HEXTAR CHEMICALS PTY. LTD ACN No.: 114 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008. Telephone: (03) 8687 9509 Fax: (03) 8687 9078

DICER 500

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and setup for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of DICER 500 Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

In a Transport
Emergency Dial
000
Police or Fire Brigade



HEXTAR CHEMICALS PTY. LTD. ACN NO.: 114 525 709

Suite 218b, Site One,
757 Bourke Street, Docklands, VIC 3008.
Telephone: (03) 8687 9509
Fax: (03) 8687 9078

Batch No.:

Date of Manufacture:



9555497505657

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DICER 500

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA present as the dimethylamine salt

GROUP I HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugarcane, turf, rice and non-crop areas as per the Directions for Use.

IMPORTANT: READ THE LABEL AND ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT

CONTENTS: 110L CONTENTS: 200L CONTENTS: 1000L

APVMA Approval No.: 65687/51582

Hextar

HEXTAR CHEMICALS PTY. LTD ACN No.: 114 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008. Telephone: (03) 8687 9509 Fax: (03) 8687 9078

DICER 500

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and setup for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of DICER 500 Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

In a Transport
Emergency Dial
000
Police or Fire Brigade



HEXTAR CHEMICALS PTY. LTD. ACN NO.: 114 525 709

Suite 218b, Site One,
757 Bourke Street, Docklands, VIC 3008.
Telephone: (03) 8687 9509
Fax: (03) 8687 9078

Batch No.:

Date of Manufacture:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DICER 500

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA
present as the dimethylamine salt

GROUP **I** HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugarcane, turf, rice and non-crop areas as per the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE USE



HEXTAR CHEMICALS PTY. LTD. (ACN No. 114 525 709)
Suite 218b, Site One, 757 Bourke Street,
Docklands, VIC 3008
Telephone: (03) 8687 9509 Fax: (03) 8687 9078

APVMA Approval No.: 65687/51582

GENERAL INSTRUCTIONS

Spray when weeds are young and actively growing, and unless stated otherwise, before the 3-5 true leaf stage.

RESISTANT WEEDS WARNING

GROUP **I** HERBICIDE

DICER 500 Selective Herbicide is a member of the Benzoic acid group of herbicides. The product has the disruptor of plant cell growth mode of action. For weed resistance management the product is a Group I Herbicide. Some naturally occurring weed biotypes resistant to the product and other Group I Herbicides may exist through normal genetic variability in any weed population. The resistant weeds can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Hextar Chemicals Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

MIXING

Add the required amount of the product directly to the tank and mix well.

APPLICATION

For Boom Application: Apply by a properly calibrated boom spray using not less than 50L of water per ha unless indicated in the Critical Comments column.

For Aerial application: Apply at least 25L total spray volume per ha. The aircraft should fly as low as practical under the prevailing conditions to minimize drift.

For High Volume Application: Apply minimum 1500L per ha.

For Knapsack Application: A 15L knapsack treats 150m².

MIMOSA CONTROL

Broadacre application: Apply 2 - 2.4 L/ha water in 60L of water per ha. Helicopter application preferred.

High Volume Application: Apply 400mL in 110L water using ground equipment. Thoroughly wet all leaves and stems of the plant.

Basal Bark Application: Apply 60mL in 15L water on plants of less than 5cm basal diameter. Spray liberally all around the base and 25 - 40cm up the stem.

Cut Stump Application: Apply 60mL in 15L water to plants up to 10cm basal diameter. Cut trees as close to ground level as possible and spray freshly cut surface before the sap dries.

Note: A temporary wilting may be evident in crops after application. If seasonal conditions are late or dry seek further advice before spraying as damage may result from spraying crops under these conditions. Growers should seek advice before spraying crops under these conditions. Growers should seek advice before spraying recently released cereal varieties. Certain maize varieties may develop clubfoot of the crop roots.

PLANT BACK PERIODS for Conservation Tillage

Wait for the following periods before planting the following crops into soils sprayed with the product. If applied to a dry soil, at least 15mm of rainfall is required before the plant back begins.

NOTE: If waterlogging occurs shortly after sowing, the product may cause some reduction in crop vigour. Use as directed only in the states indicated.

NSW, ACT, VIC, SA, WA, TAS

Rate per hectare	PLANT BACK PERIOD (days)				
	Wheat	Barley	Oats	Triticale	Rye
200mL	1	1	1	1	1
280mL	7	7	7	7	7
560mL	14	14	14	14	14

NSW, ACT, VIC, SA, WA, TAS

Rate per hectare	PLANT BACK PERIOD (days)				
	Lupins	Clover/ Medics	Mung Beans	Sunflower	Safflower
200mL	7	7	5	1	14
280mL	14	14	5	7	21
560mL	21	21	10	14	28

NSW, ACT, VIC, SA, WA, TAS

Rate per hectare	PLANT BACK PERIOD (days)			
	Pigeon Peas	Soybeans	Millet	Cotton
200mL	5	5	1	7
280mL	5	5	3	7
560mL	10	10	7	14

NSW, ACT, VIC, SA, WA, TAS

Rate per hectare	PLANT BACK PERIOD (days)				
	Sorghum	Maize	Rapeseed	Field Peas	Chick peas
200mL	1	1	7	N.D.*	N.D.*
280mL	3	3	10	14	21
560mL	7	7	14	21	28

*Not determined.

DIRECTIONS FOR USE

Restraints:

DO NOT spray when rain seems likely to occur within 4 hours.
DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or waterlogging.
DO NOT spray outside recommended crop growth as crop damage may result.
DO NOT apply to crops or pastures with clover, Lucerne or medics.

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Oats, Barley, Triticale, Commercial Rye from 5 leaf to early tillering (Zadoks 15-22)	Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Seeding Docks, Tares, Hexham Scent (Mellilotus), Wild Turnip (<i>Brassica tournefortii</i>), Soldier Thistle, Mustards, Wild radish, Capeweed, Common Ice-Plant, Charlock, Ward's Weed, Vetch, Turnip Weed (<i>Rapistrum rugosum</i>)	All states	160mL plus 700mL of MCPA 500g/L	Spray when most weeds have germinated and are in the 2 to 4 leaf stage in some instances, suppression only of Wireweed (Hogweed) and Capeweed will occur
Wheat, Oats, Barley, Triticale, Commercial Rye from 5 leaf early tillering stage until the fully tillered stage and before jointing occurs (Zadoks 15, 21-30)	Doublegee (Threecornered Jack, Spiny Emex), Wireweed, Vetch, Variegated Thistle, Tares, Sunflower, Hexham Scent (Mellilotus), Docks	SA only	280mL	Spray when most weeds have germinated and are still in the young rosette stage.
	Doublegee (Threecornered Jack, Spiny Emex), Climbing Buckwheat (Black Bindweed), Clover, Hexham Scent (Mellilotus), Docks	WA only		When weeds do not form a rosette, spray when they have not more than 8 true leave, in some instances, suppression only of Wireweed (Hogweed) and Climbing Buckwheat (Black Bindweed) will occur
	New Zealand Spinach, Climbing Buckwheat (Black Bindweed), Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Clover, Hexham Scent (Mellilotus), Docks, Fat Hen, Variegated Thistle, Scotch (Cotton or Woolly), Thistle, Sunflowers, Vetch, Tree Hogweed	QLD, NSW, ACT, VIC only		
	Wild Turnip (<i>Brassica tournefortii</i>), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron Thistle, Chickweed, Clover, Common Ice-Plant, Hoary Cress, Ward's Weed, Turnip Weed (<i>Rapistrum rugosum</i>)	SA only	280mL plus 1L of MCPA 500g/L or 560mL of 2,4-D Amine 625g/L	DO NOT use DICER 500 Selective Herbicide plus 2,4-D in oats in SA.
	Common Ice-Plant	WA only	320mL plus 640mL of 2,4-D Amine 625g/L	If any of these weeds are found in these cereal crops as well as the above weeds, the addition of MCPA or 2,4-D Amine is essential. Spray when most weeds have germinated and are still in the young rosette stage. In some instances, suppression only of Wireweed (Hogweed), Hoary Cress and Capeweed will occur
Chickweed, Sorrel, Wireweed	WA only	400mL or 280mL plus 640mL of 2,4-D Amine 625g/L or 280mL plus 1.2L of MCPA 500g/L	A mixture with 2,4-D Amine must not be used in Tasmania.	
Mintweed	NSW, ACT, QLD only	280mL plus 1.2L of MCPA 500g/L		
Wild Turnip (<i>Brassica tournefortii</i>), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron Thistle, Chickweed, Turnip Weed (<i>Rapistrum rugosum</i>)	QLD, NSW, ACT, VIC, TAS, only	280mL plus 560mL of 2,4-D Amine 625g/L		
Wheat, Barley, Triticale, Commercial Rye. From early tillering (4 to 5 fully emerged main stem leaves, plus one or more tiller) until the fully tillered stage and before the start of jointing (Zadoks 21-30)	Climbing Buckwheat (Black Bindweed), Deadnettle, Wild Turnip (<i>Brassica tournefortii</i>), Turnip Weed (<i>Rapistrum rugosum</i>)	NSW, ACT, QLD only	200-280mL plus 5-7g Metsulfuron Methyl	Spray when weeds have no more than 8 true leaves. Use higher rate on target weeds. Tank mixes of DICER 500 Selective Herbicide and Metsulfuron Methyl may result in reduced control of Wireweed in some situations
Barley, Triticale, Commercial Rye from 5 leaf until the fully tillered stage and before jointing occurs (Zadoks 21-30)	Volunteer Peas	SA only	60mL plus 1L Dicamba/MCPA	Apply when most peas are 5 - 7 cm high
Wheat, Oats from 5 leaf until the fully tillered stage and before jointing occurs (Zadoks 15-30)	Volunteer Peas	SA only	120mL plus 1L Dicamba /MCPA	Apply when most peas are 5 - 7 cm high
Rice Pre-sowing Post sowing either before permanent water or until rice is at the mid-tillering stage	Docks	NSW, ACT, NT only	400mL (seedling docks) 560mL (mature docks)	Pre-sowing: Apply before the end of August and do not cultivate or graze for 14 days after spraying. Post-sowing: Preferably apply before the application of permanent water. If permanent water has been applied DICER 500 Selective Herbicide can be applied up to the mid-tillering stage, provided the docks are exposed. Use 100L/ha for boom application.
Pinus radiata plantations At least 10 months prior to planting pines	Sorrel (in pastures to be planted to pines)	All states	1.1L or 560mL plus 1L of 2,4-D Amine 625g/L	Apply in September/October period before the sorrel flowers. Add surfactant or another non-ionic wetting agent to the mixture at label rate
	Naturally regenerated pines	SA, VIC, NSW only	2-8L	Use the high rate where total control is essential. Apply in spring. Add surfactant or another non-ionic wetting agent to the mixture at label rate.

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Sugarcane Post emergence	Bathurst Burr, Bellvine, Blackberry Nightshade, Calopo, Caltrop, Cleome, Cobbler's Pegs, Common Sow Thistle, Convolvulus, Fat Hen, Green Amaranth, Khaki Weed, Noogoora Burr, Parthenium Weed, Perennial Gooseberry, Pigweed, Pink Burr, Redroot Amaranth, Sensitive Plant, Sidaretusa, Tar Vine, Thickhead, Thornapple	NSW, ACT, QLD, WA, NT, only	560mL	Spray when most weeds have germinated and are in the young rosette stage or when they have not more than 8 leaves. Method of Application: By boom spray as an overall directed spray applying 100 to 250L water per ha or by aircraft applying 45L per ha.
	Amaranthus, Black (Giant) Pigweed, Caltrop, Mintweed, Noogoora Burr, Pigweed, Thornapple (Datura)	NSW, ACT, QLD, NT only	280mL plus 1.8L Atrazine 500g/L or 320mL plus 1.5L Atrazine 500g/L	Apply when weeds have 3 to 5 true leaves. DO NOT apply in hot, dry conditions. Apply by boom spray as an overall spray applying 50 to 250L water per ha. Use the higher rate of DICER 500 Selective Herbicide where Caltrop is present.
Potatoes	Clover, Docks, Wireweed (Hogweed), Thistles, Chickweed, Climbing Buckwheat (Black Bindweed), Black Nightshade, Fat Hen	TAS only	500mL-1.2L	Apply after haulm senescence in sufficient water to give even coverage. May be mixed with Amitrole or 2,4-D to improve spectrum of weeds controlled. Refer to respective labels of these herbicides for application rates. Add suitable surfactant for improved coverage. NOTE: Do not apply when potato haulms are still green

SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS	
			PER HA	HIGH VOL PER 100L	PER 15L KNAPSACK		
Grasses and Pastures and Perennial Grass Seed Crops	Cotton (Woolly or True Scotch) Thistle, Variegated Thistle, Star Thistle, Black or Spear Thistle, Ilyrian Thistle, Doublegee (Threecornered Jack, Spiny Emex), Parthenium Weed, Smart-Weed (Water Pepper), Hexham Scent (Mellilotus), Stemless Thistle, Caltrop (Yellow Vine), Horehound and Rumex spp. Including Curled and Broadleaf Docks	All states	280mL - 560mL for seedlings	40mL for seedlings	10mL for seedlings	When applying by boom spray use 110 to 280L of mixture per ha	
	Common Ice-Plant, Hoary Cress, Sorrel and Saffron Thistle		560mL - 1.1L for young mature plants	76mL for young mature plants	18mL for young mature plants	Add surfactant to the mixture at label rate	
			280mL plus 800mL of 2,4-D Amine 500g/L	20mL plus 60mL of 2,4-D Amine 500g/L	6mL plus 12mL of 2,4-D Amine 500g/L		
	Annual Ragweed, Blue Heliotrope, Burr Ragweed, Galvanised Burr, Ground Cherry, Ox-Eye Daisy, Perennial ragweed, Russian Knapweed, Skeleton Weed, Wild (Field or Crow) Garlic			8.8L	600mL	130mL	Use a minimum of 1500L/ha water carrier. Add surfactant. Spray Russian Knapweed at flowering. Spray Angled Onion Weed, Bladder Campion, Ox-Eye Daisy and Patterson's Curse before flowering. Spray Wild Garlic during Spring before bulbs start to form and re-spray later if dormant bulbs shoot. Spray prior to flowering.
	California (Perennial or Canada) Thistle, Field Bindweed, Patersons Curse, Prairie Ground Cherry Ragwort, Stinking Mayweed (Fetid chamomile)			4L	280mL	60mL	
	Afghan Thistle, Artichoke Thistle, Star Thistle			1.6L	100mL	24mL	
Non-crop situations	Horehound, Angled Onion Weed (Threecornered Garlic), Bladder Campion, Cotton (Woolly or True Scotch) Thistle, Docks, Sorrel, Stemless Thistle, Variegated (or Spotted) Thistle		1.2L	80mL	18mL		
	Ilyrian Thistle, Parthenium Weed, Smart-Weed		600mL	40mL	10mL		
	Poverty Weed	SA only		2L	440mL		
	Mist Flower	QLD, NT only	12L	800mL	180mL		
	Blackberry Nightshade	WA only	320 - 560mL		5 - 9mL	Blackberry Nightshade at seedling stage. For Creeping Knotweed apply in early summer to actively growing plants. For Slender Ice-Plant and Khaki Weed spray when actively growing at young rosette stage. Rubber Vine apply in autumn during active growth.	
	Creeping Knotweed		1.2 - 2.4L		18 - 36mL		
	Slender Ice-Plant		320mL plus 800mL 2,4-D 500g/L		6mL		
	Khaki Weed		600mL		9mL		
	Rubber Vine		1.4L		22mL		
	Grass Pastures and Non-Crop	Rubber Vine	QLD, NT only	2.4L plus 2.5L of 2,4-D Ester 800g/L	140 - 280mL plus 175mL 2,4-D Ester 800g/L	20 - 40mL plus 25mL of 2,4-D Ester 800g/L	Apply during April or May. For application by misting machines add 600mL DICER 500 Selective Herbicide to 10L of water

SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			PER HA	HIGH VOL PER 100L	PER 15L KNAPSACK	
Turf	Catsear, Cudweeds, Pennyworth, Lamb's Tongue, Fleabanes, Fumitory, Wireweed, Chickweed, Cotula, Sorrel, Dock, Clovers, Capeweed, Toad Rush, Peppergrass, Dandelion, Medics, Scarlet Pimpernel, Creeping Oxalis, Pearlwort, Trefoil, Yarrow, Bindy-Eye, Jo-Jo (Onehunga)	All states	1.2L plus 3.2L of 2,4-D Amine 625g/L	80mL plus 200mL of 2,4-D Amine 625g/L	18mL plus 50mL of 2,4-D Amine 625g/L	Use a minimum of 1000L/ha water. Do not spray on Buffalo Grass, Bent Grass, Lippia or Strawberry Clover. Spray when weeds are young and actively growing.
	Fat Hen, Red Flowered Mallow, Purple-Top, Stagger Weed, Swine Cress	NSW, ACT, NT, SA, QLD, WA, TAS only				
Woody Species	Moreton Bay Ash (Carbeen), Blue Gum, Bitter Bark, Brisbane Black Wattle	All states	By partial frill or tree injection technique mix 1 part DICER 500 Selective Herbicide with 12.5 parts of water and apply 2mL of the mixture to cuts in the softwood at not more than 6cm spacings around the circumference of the tree			
	Mimosa (<i>Mimosa pigra</i>)	NT only	2 - 2.4L	400mL	60mL	Apply to actively growing plants. Preferably apply during wet season. Retreatment may be necessary to control seedling growth (for application see general instructions)

ADDITIONAL RESTRAINTS FOR CONSERVATION TILLAGE

- When grass and broadleaf weeds are present, use a mixture of DICER 500 Selective Herbicide and Glyphosate 450.
- DO NOT disturb treated weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds.
- DO NOT treat weeds heavily covered with dust or silt.
- DO NOT apply if rainfall is likely within 6 hours of application.
- Observe plant back periods listed under General Instructions.
- When applying to dry soil surfaces, at least 15 mm rainfall is required before the plant back period begins.

CROP	WEEDS CONTROLLED	STATE	RATE/HA	CRITICAL COMMENTS
Seed Bed Salvage, Conservation tillage, Direct Drilling For weed control in fallow, stubble or pasture prior to sowing with a full disturbance implement	Amaranthus (Red Shank), Native Amaranth, Sow Thistle, Volunteer Cereals, Turnip Weed, Wireweed (Hogweed), Medics, Variegated Thistle, Wild Oats, Wild Mustard, Docks, Paterson's Curse	QLD, NSW, ACT, VIC, SA, WA, NT only	160 - 240mL plus glyphosate 450g/L at 400 - 600mL or glyphosate 360g/L at 500 - 700mL	If excessively wet conditions occur shortly after application or sowing some retardation of crop establishment may occur in certain broadleaf crops.
	Barnyard Grass, Liverseed Grass	NSW, ACT only		Spray when most weeds have germinated and are still in the young rosette or pre-tillering stage. Use the lower rate when weeds are prior to tillering or seedlings. Increase to the higher rate as weeds reach tillering or are 10-15 cm diameter/high. In grazed situations if heavy grazing has occurred allow re-growth to 6-8 cm high before spraying. Visible symptoms of weed control may not be apparent for 10-21 days.
	Apple of Peru, Mintweed, Noogoora Burr, Yellow Vine (Caltrop), Climbing Buckwheat (Black Bindweed), Barnyard Grass, canary Grass (Annual Phalaris), Thorn Apple, Wild or Prickly Lettuce	QLD, NSW, ACT, NT only		
	Annual Ryegrass, Volunteer Field Peas, Tree Hogweed, Capeweed	NSW, ACT, VIC, SA, WA only		
	Barnyard Grass, Liverseed Grass	QLD, NT only	160 - 240mL plus glyphosate 450g/L at 800mL - 1.2L	
	Sorrel	WA, SA, VIC, NSW, ACT only	160 - 240mL plus 650 - 800mL glyphosate 450g/L or 800mL - 1L glyphosate 360g/L	
Conservation tillage, Direct Drilling, No -till fallow, minimum tillage and all between cropping applications	Clover	All states	200mL	Add to glyphosate or SpraySeed as required (mixtures of SpraySeed plus DICER 500 Selective Herbicide must not be used in Tas). Follow label recommended rates
	Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Vetch, Docks, tares, Sunflower, Hexham Scent (Melilotus), Variegated Thistle	SA only	280mL	Spray when most weeds have germinated and are still in the young rosette stage, when they have no more than 8 true leaves
	Doublegee (Threecornered Jack, Spiny Emex), Climbing Buckwheat (Black Bindweed), Docks, Hexham Scent (Melilotus)	WA only		
	New Zealand Spinach	QLD, NSW, ACT, NT only		Spray when most weeds have germinated and are still in the young rosette stage, when they have not more than 8 true leaves
	Climbing Buckwheat (Black Bindweed), Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Hexham Scent (Melilotus), Docks, Fat Hen, Variegated Thistle, Scotch (Cotton or Woolly) Thistle, Sunflowers, Vetch, Tree Hogweed	QLD, NSW, ACT, TAS, VIC, NT only		
	Chickweed, Sorrel, Wireweed	WA only	400mL	
	Common Ice-Plant, Hoary Cress, Sorrel, Saffron Thistle	All states	280mL plus 640mL of 2,4-D Amine 625g/L	
	Caltrop (Yellow Vine), Datura, Bathurst and Noogoora Burrs, Parthenium Weed, Amaranthus, Mintweed, Annual Gooseberry, Blackberry Nightshade, fat Hen, Sunflower, Seedling Perennial Gooseberry, Seedling Khaki Weed, Bellvine, Cobbler's Pegs, Star Thistle, Black or Spear Thistle, Ilyrian Thistle, Doublegee (Threecornered Jack, Spiny Emex), Smart-Weed (Water Pepper), Stemless Thistle, Horehound and Rumex spp. Including Curled and Broadleaf Docks	All states	320 - 560mL for seedlings	Apply when weeds have at least 3 to 5 true leaves and are actively growing. Do not treat weeds, which are beyond the rosette stage. Use higher rate on large weeds.
Convolvulus, Sida-retusa, Sensitive Plant, Pink Burr, Cleome, Centro, Thickhead, Tar-Vine	QLD, NSW, NT, VIC, TAS, WA only			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

QLD, NT only

RATE	CROP	PLANT BACK PERIOD
160 - 240mL/ha	Wheat, Barley, Oats, Triticale, Commercial Rye, Sorghum, Maize, Millet, Sunflowers	1 day
	Soybeans, Mung Beans, Pigeon Peas	5 days
	Cotton/Rapeseed	7 days

NOTE:

FOR QLD, NT ONLY: FOR ALL CROPS AND RATES ABOVE 600mL/ha - PLANT BACK PERIOD 21 DAYS.

FOLLOWING POTATO HAULM DESTRUCTION IN TAS: FOR RATES AND CROPS NOT INCLUDED ABOVE, A PLANT BACK PERIOD OF 6 WEEKS APPLIES GENERALLY AND 8 WEEKS FOR SEED CROPS.

COMPATIBILITY

DICER 500 Selective Herbicide is compatible with Flowable Igran, Glyphosate 450, Glyphosate 360, Metsulfuron Methyl, Chlorsulfuron, 2,4-D amine, 2,4-D ester, Diuron, Molinate and LVE MCPA. Also compatible with Atrazine, but when used in combination with this product some temporary visible effects in the crop growth may be observed.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. To prevent damage, DO NOT permit drift or vapour mist to come into contact with fruit and ornamental trees, vines, vegetables, flowers, tobacco or any other field crop. Minimise spray drift by using low pressure and nozzles, which DO NOT give a fine droplet size. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use in high wind. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT use at higher rate than recommended.

PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT

DO NOT use container for any other purposes
DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site, if recycling, replace and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and setup for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of DICER 500 Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

In a Transport
Emergency Dial
000
Police or Fire Brigade



HEXTAR CHEMICALS PTY. LTD. (ACN No: 114 525 709)
Suite 218b, Site One, 757 Bourke Street,
Docklands, VIC 3008
Telephone: (03) 8687 9509 Fax: (03) 8687 9078

APVMA Approval No. 65687/51582

Name of File:
65687_51582_leaflet_v1.pdf

Dimensions:
36 cm (W) x 46.2 cm (H)

Scale of Print:
A4 55%
A3 80%