

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

WELOXY 520 EC HERBICIDE

ACTIVE CONSTITUENT: 520g/L HALOXYFOP
present as the haloxyfop-R-methyl ester

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions for Use.

IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT

CONTENTS: 5 Litres

APVMA Approval No.: 66570/53913

Hextar

HEXTAR CHEMICALS PTY LTD ABN 11 134 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008. Telephone: (03) 8687 9509 Fax: (03) 8687 9078

WELOXY 520 EC HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near feedstuffs, fertilisers or seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of Weloxyl 520 EC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.



IN A TRANSPORT
EMERGENCY
DIAL 000
Police or Fire Brigade

Batch No.:

Date of Manufacture:



9555497505978

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

WELOXY 520 EC HERBICIDE

ACTIVE CONSTITUENT: 520g/L HALOXYFOP
present as the haloxyfop-R-methyl ester

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT**

CONTENTS: 10 Litres

APVMA Approval No.: 66570/53913

Hextar

HEXTAR CHEMICALS PTY LTD ABN 11 114 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008. Telephone: (03) 8687 9509 Fax: (03) 8687 9078

WELOXY 520 EC HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near feedstuffs, fertilisers or seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of Weloxyl 520 EC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.



IN A TRANSPORT
EMERGENCY
DIAL 000
Police or Fire Brigade

Batch No.:

Date of Manufacture:



9555497505985

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

WELOXY 520 EC HERBICIDE

ACTIVE CONSTITUENT: 520g/L HALOXYFOP
present as the haloxyfop-R-methyl ester

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT**

CONTENTS: 20 Litres

APVMA Approval No.: 66570/53913

Hextar

HEXTAR CHEMICALS PTY LTD ABN 11 114 525 709

Suite 218b, Site One, 757 Bourke Street, Docklands, VIC 3008. Telephone: (03) 8687 9509 Fax: (03) 8687 9078

WELOXY 520 EC HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near feedstuffs, fertilisers or seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of Weloxy 520 EC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.



IN A TRANSPORT
EMERGENCY
DIAL 000
Police or Fire Brigade

Batch No.:

Date of Manufacture:



9555497505992

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

WELOXY 520 EC HERBICIDE

ACTIVE CONSTITUENT: 520g/L HALOXYFOP
present as the haloxyfop-R-methyl ester

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT



HEXTAR CHEMICALS PTY. LTD. (ABN 11 114 525 709)
Suite 218b, Site One, 757 Bourke Street,
Docklands, VIC 3008.
Telephone: (03) 8687 9509 Fax: (03) 8687 9078

APVMA Approval No.: 66570/53913

DIRECTIONS FOR USE

Restraints:
DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result.
DO NOT spray if rain is likely to occur within one hour.

Table 1a. Winter crops – Canola, Chickpeas, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Lucerne, Vetch, Medic and Clover pastures or seed crops

Weeds Controlled	Weed Growth Stage	Rate (mL/ha)		Critical Comments
		With Uptake Spraying Oil ¹	With non-ionic wetter ²	
				^{1,2} See GENERAL INSTRUCTIONS , Spraying oils/ wetters section.
Annual Ryegrass	2 to 4 leaf	75	100	FIELD PEAS AND CANOLA: The only oil recommended for use with Weloxy 520 is Uptake* Spraying Oil. Weloxy 520 + Corporal 750 SG + Uptake* Spraying Oil are compatible and selective to canola. This tank-mix is also compatible with atrazine and selective to triazine tolerant canola.
	early tillering	100	100	
Barley Grass Brome grass Paradoxa grass Volunteer cereals	2 to 4 leaf	50	75	LUPINS AND FIELD PEAS: Mixtures with diflufenican or simazine may cause crop yellowing and separate applications are recommended.
	early tillering	75	100	
Wild Oats WA, SA, Vic, Tas, southern & central NSW	2 to 4 leaf	37.5	50	CHICKPEAS, FABA BEANS, LENTILS AND VETCH, LINOLA, LINSEED: Broadleaf herbicides should not be added to Weloxy 520. Apply Weloxy 520 and broadleaf herbicides at least a week apart.
	early tillering	50	75	
Wild Oats northern NSW & Qld	2 to 4 leaf	50	75	LUCERNE, CLOVER OR MEDIC PASTURES: If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS , Spraying oils/wetter section). If silver grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank mixed with the higher rate of Weloxy 520 plus a non-ionic wetter.
	early tillering	75	100	

Table 1b. Winter crop growth stage application windows

Crop	Crop Growth Stage
Lucerne, Medic and Clover pastures or seed crops.	Apply from 2 nd trifoliolate leaf onwards. For <i>Erodium</i> spp spraying, apply from cotyledon crop stage onwards.
Canola	Apply from 2 nd leaf and prior to bud formation and stem elongation.
Linola, Linseed	Apply from 5cm to flowering.
Chickpeas, Faba beans, Field peas, Lentils, Lupins, Vetch	Apply from 2 nd leaf, 2 nd node or 2 nd branch prior to flowering.

Table 2a. Lucerne, Medic and Clover seed crops and pastures. See table 1b for crop stages.

Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake Spraying Oil	Critical Comments
Prairie grass (<i>Bromus catharticus</i>)	Up to early tillering	100	¹ See GENERAL INSTRUCTIONS , Spraying oils/wetters section.
Musky or ferny leaf Storksbill: (<i>Erodium moschatum</i>) Common Crowsfoot or Common Storksbill (<i>Erodium cicutarium</i>)	Up to 6 leaf or 5cm diameter	50 – 75 ³	³ Use lower when growing conditions and crop or pastures competition are good and when weed populations are below 100 plants/m ² . Use the higher rate when weed population re above 100 plants/m ² or when crop or pasture competition is poor. NOTE: Storksbill may not be controlled if simazine or flumetsulam are tank-mixed with Weloxy 520 EC Herbicide.
Long or shiny leaf Storksbill (<i>E. botrys</i>)	Up to 8 leaf or 5cm diameter	75 - 100	LUCERNE, CLOVER OR MEDIC PASTURES If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75mL + spraying oil or 100mL + wetter/ha. (See GENERAL INSTRUCTIONS , Spraying oils/wetter section). If Silver grass (<i>Vulpia</i> spp.) is presented in pasture, simazine should be tank mixed with the higher rate of Weloxy 520 EC Herbicide plus a non-ionic wetter.

Table 2b. Lucerne, Medic and Clover seed crops only – not to be used for stockfeed. See table 1b for crop stages

Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake Spraying Oil	Critical Comments
Couch grass (suppression), Rhodes grass (control)	Tillering seedlings	150 + 150 ⁴	⁴ See GENERAL INSTRUCTIONS , Spraying oils/wetters section.
Couch grass (control), Rhodes grass (control)	Established stands	400 – 800	⁴ For best suppression of couch or control of Rhodes grass, make 2 applications of Weloxy 520 EC Herbicide 2-4 weeks apart. Time second application to coincide with tillering stage of weeds and just after irrigation or significant rain. Only treat actively growing weeds, which are not moisture stressed. Use these rates for control of couch and Rhodes grass.

Table 3a. Summer Crops – Cotton, Cowpea, Lucerne, Mung bean, Navy beans, Peanuts, Soybeans, Sunflowers

Weeds Controlled	Weed Growth Stage	Rate (mL/ha) with Uptake Spraying Oil	Critical Comments
Australian millet	2 leaf to tillering up to 15cm	150	¹ See GENERAL INSTRUCTIONS , Spraying oils/wetters section.
Barnyard grass	2 to 5 leaf	100	NAVY BEAN, PEANUTS, SOYBEANS For broadleaf weed control, Weloxy 520 EC Herbicide at 150mL/ha plus wetter may be tank mixed with Acifluorfen (except on Navy beans) or Bentazone. Tank mixes may cause transient leaf spotting on the crop but do not normally affect yield. DO NOT tank mix broadleaf herbicides with Weloxy 520 EC Herbicide if grasses have begun tillering or if the grasses are under moisture stress. DO NOT add Uptake Spraying Oil with Acifluorfen or Bentazone. DO NOT use Acifluorfen or Bentazone tank-mixes on Cowpeas.
	tillering up to 15cm	150	
Crowsfoot grass Green Panic grass Johnson grass (Rhizome)	2 leaf to tillering up to 15cm	150	
Johnson grass (seedling) Liverseed grass (seedling) Mossman River grass	2 to 5 leaf	100	
	tillering up to 15cm	150	
Summer grass	2 leaf to tillering up to 15cm	150	
Volunteer cereals	2 to 4 leaf	100	
	tillering up to 15cm	150	

Table 3b. Summer crop growth stage application windows

Crop	Crop Growth Stage
Lucerne	Apply from 2 nd trifoliolate leaf onwards
Cowpea, Mung beans, Navy beans, Soybeans	Apply from 2 nd leaf to flowering
Peanuts	Apply from 5cm to pegging
Cotton	Apply from 2 nd leaf to before the onset of flowering
Sunflowers	Apply from 2 nd leaf to head initiation

GENERAL INSTRUCTIONS MIXING

- Add water to the spray tank to 10cm above the level of agitation and ensure the agitation device is working vigorously (There must be a minimum of 100L of water in the tank before any pesticide is added).
- If tank mixing, firstly add any soluble liquid formulations (e.g. Clopyralid 300 AC Herbicide) and allow agitation for approximately 1 minute.
- Then add Weloxy 520 EC Herbicide at the pot where the agitation is the strongest (do not add Weloxy 520 EC Herbicide through a strainer or sieve). Allow further agitation for 1 minute.
- Half fill spray tank.
- If using wettable powder or water dispersible granules, or other emulsifiable concentration formulations (e.g. Chlorpyrifos 750 WG or Omethoate), these should be added after the Weloxy 520 EC Herbicide to the half full spray tank to ensure vigorous agitation.
- Finally add Uptake Spraying Oil or approved alternate spraying oil/wetter (see section on spraying oils/wetters) and continue filling the tank to the required volume maintaining agitation at all times
- Only mix sufficient solution for immediate use. Weloxy 520 EC Herbicide and any other tank mixes should be applied immediately for best results.

Spraying Oils/Wetters

***Spraying Oils:** It is essential to add an adjuvant to Weloxy 520 EC Herbicide. Best results will be achieved with Uptake Spraying Oil at 0.5L/100L of spray solution. Alternatively, other oils plus non-ionic wetter may be used. When other crop spraying oils are used, mix at 1.0L/100L and add a non-ionic wetter (surfactant) at 200mL/100L of spray solution. Use of an oil is not always recommended. See Critical Comments for specific situation recommendations.
†Non-ionic Wetters: When Uptake or other oils are not used, a 100% concentrate non-ionic wetting agent such as BS-1000 at 200mL/100L must be used along with the higher rate of Weloxy 520 EC Herbicide as specified in the Directions for Use. Where water volumes of less than 50L/ha are used, DO NOT use less than 250mL/ha of Uptake or 500mL/ha for oils other than Uptake or less than 100mL of wetter.

CANOLA, LUCERNE, MEDIC AND CLOVER PASTURES AND SEED CROPS

When tank mixing Weloxy 520 EC Herbicide with Clopyralid 300 AC herbicides (canola only) or Flumetsulam (lucerne, clover and medics), use Uptake Spraying Oil with lower rates of Weloxy 520 EC Herbicide or a wetting agent with the higher rates of Weloxy 520 EC Herbicide unless otherwise specified. When mixing Weloxy 520 EC Herbicide with other broadleaf herbicides on these crops, DO NOT use an oil use a wetter instead.

FIELD PEAS AND CANOLA

The oil recommended is Uptake Spraying Oil. Hasten is also recommended for use with tank-mixtures of Weloxy 520 EC Herbicide and Clethodim 240 EC Herbicide.
For canola, Weloxy 520 EC Herbicide + Corporal 750SG + Uptake Spraying Oil are compatible and selective to canola. This tank-mixture is also compatible with atrazine or simazine and selective to triazine tolerant canola.

NAVY BEANS, PEANUTS AND SOYBEANS

When mixing with Acifluorfen or Bentazone DO NOT add spraying oil to these mixtures. DO NOT use these tank-mixtures on cowpea.

COMPATIBILITY - Ground Use Only

Insecticides: dimethoate, endosulfan, Chemicide 500 Insecticide, chlorpyrifos 750WG, omethoate

Herbicides: atrazine, bentazone, acifluorfen, flumetsulam, clopyralid 300AC, Corporal 750SG, MCPA Ester (LVE) – (Do not exceed 700mL/ha of MCPA LVE), oryzalin, clethodim 240EC, simazine, fluroxypyr 200EC

Fungicides: mancozeb DF, Mancozz 750 WG Fungicide.

Trace elements: magnesium sulphate, zinc sulphate
Weloxy 520 EC Herbicide is not compatible with 2,4-D or MPA as sodium or amine salts.

Aerial Use

No product other than a recommended crop oil or wetter should be mixed with Weloxy 520 EC Herbicide when applied by air except for the addition of Clopyralid 750 SG Herbicide for use in forestry and Corporal 750 SG Herbicide for use in canola only.

APPLICATION

Apply Weloxy 520 EC Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft delivering droplets with a VMD of 200-300 microns. The following spray volumes are recommended;
Ground Application: 50-150 L/ha
Aerial application: 30 L/ha minimum
Use higher water volumes in orchards and in dense crops where the weeds may be shielded by the crop canopy.

CLEANING SPRAY EQUIPMENT

Before using Weloxy 520 EC Herbicide
If broadleaf herbicides, particularly sulfonylureas, have been used in the spray equipment at any time prior to the use of Weloxy 520 EC Herbicide. Particular care should be taken to follow the directions on the relevant broadleaf herbicide label for equipment cleaning, or damage to susceptible crops may occur.

After using Weloxy 520 EC Herbicide

Empty the tank and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean filters in the tank, pump line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate: Before spraying cereals, maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. SURF, Cold Water SURF Concentrate, DynamMatic Concentrate, OMO or Drive) at 500mL/100L of water or the powder equivalent at 500g/100L of water, and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean separately. Finally flush the system with clean water and allow to drain. Chlorine based cleaners are not recommended.
Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

Table 4. Annual and Perennial grasses and Erodium spp. In Orchard, Vine, and Plantation crops, Forestry and Pyrethrum

Crop	Crop Growth Stage	Weeds Controlled	Weed Stage	Rate (mL/ha) with Uptake ¹ Spraying Oil	Critical Comments
Orchard, Vine and plantation Crops including: Apples, Avocado, Banana, Blueberry, Citrus, Custard apple, Feijoa, Grapevines, Guava, Kiwifruit, Litchi (Lychee), Longan, Mango, Nut trees, Passionfruit, Paw paw, Pear, Persimmon, Pineapple, Rambutan, Stone Fruit	All growth stages	Perennial grasses, Couch, Rhodes grass, Slender Rats Tail grass	Established stands	400 - 800	¹ See GENERAL INSTRUCTIONS , Spraying oils/wetters section.
		Buffel grass, Green panic, Johnson grass, Kikuyu, <i>Paspalum</i> spp, <i>Setaria</i> spp	Vegetative to early tillering	200	Spray should be directed to the base of the tree or vine avoiding contact with frit and foliage.
			Late tillering	400	
		Annual grasses: Annual ryegrass, Barley grass, Barnyard grass, Brome grass, Crowsfoot grass, Lesser Canary grass, Liverseed grass, Mossman river grass, Paradoxa grass, Summer grass, Volunteer cereals, Wild Oats	2 leaf to tillering	200	Spot spray: Use 25mL to 50mL/100L of water. Use higher rate on late tillering mature grasses. Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled.
Forestry: <i>Pinus radiata</i> , <i>Eucalyptus</i> spp.		Annual grasses as above	Vegetative to tillering	125 - 250	Forestry: For annual grasses apply lowest rate to newly emerged grasses, increasing the rate as they develop.
Forestry: <i>Pinus pineaster</i>		Annual grasses as above	Vegetative to tillering	100 - 250	Pyrethrum (Tasmania only) For <i>Erodium</i> spp apply 75-100 mL/ha if the main weed is <i>E. botrys</i> . Use 50-75 mL/ha if either <i>E. cicutarium</i> or <i>E. moschatum</i> are the main weeds.

Table 5. Weloxy 520 EC Herbicide and Clethodim 240 EC Herbicide tank mixes – Canola, Chickpeas, Faba beans, Field peas, Lupins, Lentils

Weeds Controlled	Weed Growth Stage	Rate (mL/ha)		Critical Comments
		Weloxy 520 EC Herbicide	Clethodim 240 EC Herbicide	
FOP/DIM susceptible Annual ryegrass + Volunteer barley, Volunteer wheat, Brome grass, Wild Oats, Barley grass, Phalaris	2 to 4 leaf	25	150	¹ See GENERAL INSTRUCTIONS , Spraying oils/wetters section.
	Early tillering	38	150	Use Uptake Spraying Oil at 500mL/100L or Hasten at 1L/100L.
FOP resistant Annual ryegrass + Volunteer barley, Volunteer wheat, Brome grass, Wild Oats, Barley grass, Phalaris	2 to 4 leaf	25	200	Apply at the same crop growth stages as those in Table 1b Winter Crops.
	Early tillering	38	250	Lentils: Apply up to 7 node-early branching crop growth stage. Lupins: Not for Qld.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

HARVESTING WITHHOLDING PERIODS

Canola, Chickpeas, Cotton, Cowpea, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers and Vetch:
NOT REQUIRED WHEN USED AS DIRECTED.

Medic and Clover seed crops:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

STOCK FOOD WITHHOLDING PERIODS

Canola, Chickpeas, Cotton, Cowpea, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers and Vetch:
DO NOT GRAZE OR CUT FOR STOCK FOR 28 DAYS AFTER APPLICATION

Lucerne: **DO NOT GRAZE OR CUT FOR 21 DAYS AFTER APPLICATION**

Medic and Clover pasture: **DO NOT GRAZE OR CUT FOR 7 DAYS AFTER APPLICATION**

COTTON GIN TRASH MUST NOT BE FED TO ANIMALS

RESISTANT WEEDS WARNING

GROUP A HERBICIDE

Weloxy 520 EC Herbicide is a member of the cyclohexanedione group of herbicides. Weloxy 520 EC Herbicide has the inhibition of acetyl CoA carboxylase mode of action. For weed resistance management, Weloxy 520 EC Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to Weloxy 520 EC Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Weloxy 520 EC Herbicide or other Group A herbicides.
Since the occurrence of resistant weeds is difficult to detect prior to use, Hextar Chemicals Pty Ltd accepts no liability for any losses that may result from the failure of Weloxy 520 EC Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Weloxy 520 EC Herbicide damages cereals and grasses. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible crops/plants, cropping lands or pastures. Cereal crops or grasses planted within twelve weeks of application may be damaged by the residual effects of Weloxy 520 EC Herbicide Herbicide, particularly on light and red soils.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Weloxy 520 EC Herbicide is toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near feedstuffs, fertilisers or seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

STORAGE AND DISPOSAL (cont.)

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain n a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

The use of Weloxy 520 EC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Hextar Chemicals Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Hextar Chemicals Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

In a Transport
Emergency Dial
000
Police or Fire Brigade